

ALMOST EVERY INTERVAL TRANSLATION MAP OF THREE INTERVALS IS FINITE TYPE

D. VOLK

ABSTRACT. Interval translation maps (ITMs) are non-invertible generalizations of interval exchanges (IETs). The dynamics of finite type ITMs is similar to IETs, while infinite type ITMs are known to exhibit new interesting effects. In this paper, we prove the finiteness conjecture for the ITMs of three intervals. Namely, the subset of ITMs of finite type is open, dense, and has full Lebesgue measure. For this, we show that any ITM of three intervals can be reduced either to a rotation or to a double rotation.

1. INTRODUCTION AND MAIN RESULT

1.1. Interval translation maps. Let $\Omega \subset \mathbb{R}$ be a semi-interval split into d disjoint semi-intervals, $\Omega = \sqcup_{j=1}^d \Delta_j$, $\Delta_j = [\beta_{j-1}, \beta_j)$. An *interval translation* $T: \Omega \rightarrow \Omega$ is a map given by a translation on each of Δ_j , $T|_{\Delta_j}: x \mapsto x + \gamma_j$, the vector $(\gamma_1, \dots, \gamma_d)$ is fixed. After normalization $\Omega := [0, 1)$, the space $\text{ITM}(d)$ of d intervals' translations is a convex polytope in \mathbb{R}^{2d-1} . We endow it with the Euclidean metric and the Lebesgue measure. An example of an ITM is shown on Figure 1.1. We draw Ω as a horizontal line, the splitting of Ω into Δ_j as arcs of different styles atop of the line, and the images of Δ_j as the mirrored arcs below the line.



FIGURE 1.1. An interval translation map of 3 intervals.

Interval translation maps (ITMs) were first introduced in 1995 by Boshernitzan, Kornfeld in [1]. They are a generalization of interval exchange transformations. Unlike IETs the ITMs are generally not invertible.

Define $\Omega_0 = \Omega$, $\Omega_n = T\Omega_{n-1}$ for $n \geq 1$, and let X be the closure of $\bigcap_{n=1}^{\infty} \Omega_n$. An interval translation map is called of *finite type* if $\Omega_{n+1} = \Omega_n$ for some n , otherwise it is called of *infinite type*. Denote the set of infinite type ITMs by \mathcal{S} .

2010 *Mathematics Subject Classification.* Primary: 37C05, 37C20, 37C70, 37D20, 37D45.

Key words and phrases. Dynamical systems, attractors, interval translation maps, interval exchange maps, double rotations, color rotations.

The author was supported in part by grants RFBR 12-01-31241-moLa, President's of Russia MK-2790.2011.1, "Young SISSA Scientists", and NSF IIS-1018433 (PI Max Welling, Co-PI Anton Gorodetski).

Schmeling and Troubetzkoy [10] proved that for any d , an ITM is of finite type iff X consists of a finite union of intervals. In this case, some power of the restriction $T|_X$ is an interval exchange transformation. If T is of infinite type and $T|_X$ is transitive, then X is a Cantor set. Additionally, Bruin and Troubetzkoy [4] showed that if X is transitive, then it is minimal i.e. every orbit is dense.

In their pioneering paper [1], Boshernitzan and Kornfeld gave the first example of an ITM of infinite type. They also asked, “*To what extent the ITMs of finite type are typical? In particular, is it true that ‘almost all’ ITMs are of finite type?*” Then they remarked that *the answer is affirmative for $d = 3$ but gave no proof of that.*

Until now, the question was answered (positively) only for some specific families of ITMs: by Bruin, Troubetzkoy [4] for a 2-parameter family in $\text{ITM}(3)$, by Bruin [2] for a d -parameter family in $\text{ITM}(d + 1)$, and by Suzuki, Ito, Aihara [12] and Bruin, Clack [3] for so-called double rotations, see Subsection 1.3. Schmeling, Troubetzkoy [10] established a number of related topological results, in particular, that the interval translation mappings of infinite type form a G_δ subset of the set of all interval translation mappings, whereas the interval translation mappings of finite type contain an open subset.

But the original question stood open. In this paper, we finally justify the affirmative answer for general ITMs of three intervals:

Theorem 1.1. *In the space $\text{ITM}(3)$, the set $\mathcal{S} \cap \text{ITM}(3)$ has zero Lebesgue measure.*

Remark 1.2. The numerical estimate by [3] infers that its Hausdorff dimension H satisfies $4 \leq H \leq 4.88$.

The same question about $d > 3$ intervals remains open.

The entropy and word growth properties of piecewise translations, including their higher dimension generalizations, was studied in [7], [8], [5], [9]. The question of their (unique) ergodicity was considered by [4], [6], [3].

We always assume $\gamma_i \neq 0$ and $\beta_i - \beta_{i-1} \neq 0$ for all $1 \leq i \leq d$.

1.2. Tight ITMs. Let T be any interval translation map. We say that an interval $\Delta \subset \Omega$ is *T -regular* (or simply *regular*) if there exists $N \in \mathbb{N}$ such that for any $x \in [0, 1)$ there exists $1 \leq n < N$ such that $T^n x \in \Delta$. In particular, every point of Δ returns to Δ after a uniformly bounded number of iterates of T , so the *induced* (i.e. *first-return*) map of T is well defined on Δ . We denote the induced map by T_Δ . We say that an interval $\Delta \subset \Omega$ is a *trap* if it is regular and $T\Delta \subset \Delta$. In this case, we have $T_\Delta = T|_\Delta$.

The following Lemma shows that the properties to be finite or infinite type are preserved by inductions.

Lemma 1.3. *Assume $T|_X$ is transitive. Then,*

- i) if there exists a regular Δ such that T_Δ has finite type, then T has finite type;
- ii) if T is finite type, then for any regular Δ the map T_Δ has finite type.

Proof. First we prove **i**. Assume T has infinite type. Then X is a T -invariant Cantor set with minimal dynamics (see [10], [4]). Because Δ is regular, $Y = X \cap \Delta$ is a nonempty Cantor set. It is also T_Δ -invariant, and thus $Y \subset X(T_\Delta)$. Because T_Δ has finite type, $X(T_\Delta)$ is a finite union of intervals with an IET on them. This implies $Y \neq X(T_\Delta)$. Because X is transitive, it is minimal for T , and thus $X(T_\Delta)$ is minimal for T_Δ . But we have just shown that it contains another invariant closed set. This contradiction proves the lemma.

Recall that X is a finite union of intervals with an IET on them. So, because X is transitive, it is also minimal.

Now we prove **ii**. Assume T_Δ has infinite type. Then $X(T_\Delta)$ is a T_Δ -invariant transitive set. Denote by Y the union of its N iterates by T :

$$Y = \bigcup_{n=1}^N T^n X(T_\Delta).$$

Now Y is a T -invariant closed set and $Y \subsetneq X$. This contradicts the minimality of X and thus proves the lemma. \square

For any $M \subset [0, 1]$, we denote $[M] := [\inf M, \sup M]$. We say that an interval translation map $T: \Omega \rightarrow \Omega$ is *tight* if $[T\Omega] = [\Omega]$. We denote the space of tight interval translation maps of d intervals by $\text{TITM}(d)$. Recall that $\text{ITM}(d)$ is a filled convex polytope in \mathbb{R}^{2d-1} , and observe that $\text{TITM}(d)$ is a connected finite union of filled convex polytopes in \mathbb{R}^{2d-3} .

It is easy to see that for any $T \in \text{ITM}(d)$, there exists a trap Δ such that the map T_Δ is a tight interval translation map of r intervals, $r \leq d$. Namely, take $X'_0 := \Omega$, and let $X'_k := [TX'_{k-1}]$, $\Delta := \bigcap_{k=0}^\infty X'_k$. Then $\Delta \supset X$, Δ is a trap, and T_Δ is tight.

However, to prove Theorem 1.1, we will need more explicit way to produce such a Δ .

For $d = 2$, we have $\Delta = [\beta_1 + \gamma_2, \beta_1 + \gamma_1]$, see Figure 1.2 and keep in mind that $\gamma_1 > 0$, $\gamma_2 < 0$. Indeed, for any $x < \beta_1$ we have $Tx = x + \gamma_1$. Thus there exists $1 \leq n_1 \leq \left\lceil \frac{\beta_1}{\gamma_1} \right\rceil + 1$




FIGURE 1.2. ITM of two intervals: rotation.

such that $T^{n_1}x \in [\beta_1, \beta_1 + \gamma_1)$. Similarly, for any $x \geq \beta_1$ there exists $1 \leq n_2 \leq \left\lceil \frac{1-\beta_1}{|\gamma_2|} \right\rceil + 1$ such that $T^{n_2}x \in [\beta_1 + \gamma_2, \beta_1)$. Thus Δ is regular. Now note that $T|_{\Delta}$ is just a rotation $x \mapsto x + \gamma_1 \pmod{\Delta}$, so $T\Delta = \Delta$. Thus Δ is a trap and T_{Δ} is tight.

To obtain a similar formula for $d \geq 3$, consider two cases:

$$(1.1) \quad \Delta_1 \cap T\Omega = \emptyset \text{ or } \Delta_d \cap T\Omega = \emptyset;$$

$$(1.2) \quad \Delta_1 \cap T\Omega \neq \emptyset \text{ and } \Delta_d \cap T\Omega \neq \emptyset.$$

In the first case, we can completely remove Δ_1 or Δ_d and thus reduce the problem to the study of $\text{ITM}(d-1)$. For $\text{ITM}(3)$, this case can be ignored because then the map is necessary a rotation, and thus finite type.

In the second case, let I_- be the set $\{i \mid \gamma_i < 0\}$ and I_+ be the set $\{i \mid \gamma_i > 0\}$. Because $\gamma_1 > 0$ and $\gamma_d < 0$, the both sets are nonempty. Take the interval

$$\Delta = [\delta_0, \delta_1) = \left[\min_{i \in I_-} (\beta_{i-1} + \gamma_i), \max_{i \in I_+} (\beta_i + \gamma_i) \right),$$

see Figure 1.3. In the same way as for $d = 2$ one can show that for any $x < \delta_0$ or $x > \delta_1$




FIGURE 1.3. Fitting operator.

there exists a bounded n such that $T^n x \in \Delta$. Now, for any $x \in \Delta$, x either moves left or right. Assume it moves left. Then $\min_{i \in I_-} (\beta_{i-1} + \gamma_i) \leq Tx < x$ which implies $Tx \in \Delta$. Similarly, for x moving right we have $x < Tx \leq \max_{i \in I_+} (\beta_i + \gamma_i)$. Thus $Tx \in \Delta$, and Δ is a trap.

Because we are in the second case, for every $2 \leq i \leq d$, the left end β_{i-1} of Δ_i belongs to Δ . In particular, this is true for the $i = \operatorname{argmin}_{i \in I_-} (\beta_{i-1} + \gamma_i)$. So the left end of Δ_i maps to δ_0 . Thus T_{Δ} is tight from the left. Similarly, we show T_{Δ} is tight from the right and thus tight in general.

Rescale the map T_{Δ} so that Δ becomes $[0, 1)$. We say that the result, \tilde{T}_{Δ} , is the *fitting* of T and denote the *fitting operator* $T \mapsto \tilde{T}_{\Delta}$ by $\mathcal{F}_d: \text{ITM}(d) \rightarrow \text{TITM}(d)$. When the value of d is clear, we will write \mathcal{F} instead of \mathcal{F}_d for brevity.

Proposition 1.4. *In the case (1.2), the fitting operator is a piecewise rational map of the maximal rank $2d - 3$.*

Proof. We partition the space $\text{ITM}(d)$ into the union of the cells C_{jk} , $j \neq k$:

$$C_{jk} = \{T \in \text{ITM}(d) \mid j = \operatorname{argmin}_{i \in I_-} (\beta_{i-1} + \gamma_i), k = \operatorname{argmax}_{i \in I_+} (\beta_i + \gamma_i)\}.$$

The fitting operator is the composition of truncation and rescaling:

$$\mathcal{F} = \mathcal{R} \circ \mathcal{T}.$$

The rescaling part is a rational map $\mathcal{R}: \mathbb{R}^{2d-1} \rightarrow \mathbb{R}^{2d-3}$ of rank $2d-3$. To understand the truncation part \mathcal{T} , we introduce new coordinates $B_{i-1} = \beta_{i-1} + \gamma_i$, $i = 1, \dots, d$, which are the images of the left ends of Δ_i . In particular, $B_0 = \gamma_0$. At every cell C_{jk} , the truncation is a linear map $\mathcal{T}: \mathbb{R}^{2d-1} \rightarrow \mathbb{R}^{2d-1}$. Let us show it is invertible. In the coordinates (β_i, B_i) , the truncation has the form

$$(1.3) \quad \begin{array}{c} \left[\begin{array}{c} 0, \beta_1, \beta_2, \dots, \beta_{d-2}, \beta_{d-1}, 1 \\ B_0, B_1, \dots, B_{j-1}, \dots, B_k, \dots, B_{d-2}, B_{d-1} \end{array} \right] \\ \downarrow \mathcal{T} \\ \left[\begin{array}{c} B_{j-1}, \beta_1, \beta_2, \dots, \beta_{d-2}, \beta_{d-1}, B_k \\ B_{j-1} + B_0, B_1, \dots, B_{j-1}, \dots, B_k, \dots, B_{d-2}, B_k + B_{d-1} - \beta_{d-1} \end{array} \right] \end{array}$$

Note that most of the coordinates are mapped identically. The non-identical part of (1.3) is

$$(1.4) \quad \begin{array}{c} \left[\begin{array}{c} 0, \beta_{d-1}, 1 \\ B_0, B_{j-1}, B_k, B_{d-1} \end{array} \right] \\ \downarrow \mathcal{T} \\ \left[\begin{array}{c} B_{j-1}, \beta_{d-1}, B_k \\ B_{j-1} + B_0, B_{j-1}, B_k, B_k + B_{d-1} - \beta_{d-1} \end{array} \right] \end{array}$$

which is clearly invertible. So, at every cell C_{jk} the fitting \mathcal{F} is a rational map of rank $2d-3$. \square

All the previous considerations are valid for any number of intervals. In what follows, we have the proofs only for the case of $d = 3$.

1.3. Double rotations. Following Suzuki, Ito, Aihara [12], we introduce the family of double rotations. A *double rotation* is a map $f_{(a,b,c)}: [0, 1) \rightarrow [0, 1)$ defined by

$$f_{(a,b,c)}(x) = \begin{cases} \{x + a\}, & \text{if } x \in [0, c), \\ \{x + b\}, & \text{if } x \in [c, 1). \end{cases}$$

In the circle representation $[0, 1) \equiv S^1$, a double rotation is a map $S^1 \rightarrow S^1$ defined by independent rotations of two complementary arcs of S^1 . Clearly, any double rotation is an ITM of two to four intervals. Denote the space of parameters (a, b, c) of double rotations by $\text{Rot}(2) = [0, 1) \times [0, 1) \times [0, 1)$. In the same way we define the family $\text{Rot}(n)$ of *n-rotations*.

For the double rotations, the finiteness problem is solved by Bruin and Clack in [3] using the renormalization operator of Suzuki, Ito, Aihara [12]. Namely, they proved the following

Theorem 1.5 (Bruin, Clack, 2012). *In the space $\text{Rot}(2)$, the set $\mathcal{R} = \mathcal{S} \cap \text{Rot}(2)$ of ITMs of infinite type has zero Lebesgue measure.*

In the present paper we show that the finiteness problem for ITM(3) reduces to the one for the double rotations.

Theorem 1.6. *The space TITM(3) splits into countably many open sets A, A', B, B_i, C_i , $i \in \mathbb{N}$, such that their union U is dense in TITM(3), and the complement $K = \text{TITM}(3) \setminus U$ which is a union of countably many hyperplanes. Moreover,*

- *any $T \in A \cup A'$ is a double rotation,*
- *any $T \in B$ is reduced to a double rotation via a single Type 1 induction,*
- *for any $i \in \mathbb{N}$, any $T \in B_i$ is reduced to a double rotation via a single Type 2 induction.*
- *for any $i \in \mathbb{N}$, any $T \in C_i$ is reduced to a single rotation via a single Type 2 induction.*

On every piece A, A', B, B_i, C_i , these inductions are invertible rational maps.

Conjecture 1.7. *The space TITM(d) splits into countably many open sets A_i such that their union U is dense in TITM(d), and the complement $K_d = \text{TITM}(d) \setminus U$ which is a union of countably many hyperplanes. Any $T \in U$ is reduced to an $(d - 1)$ -rotation via some induction.*

Because the inductions are local diffeomorphisms, the preimages of zero measure sets are zero measure sets. The Hausdorff dimension is also preserved. This allows to transfer the results of [12] and [3] to TITM(3).

2. PROOF OF THEOREM 1.6

Let $T \in \text{TITM}(3)$ with $\Omega = [0, 1)$. Without loss of generality, we can assume $\gamma_1 > 0$ and $\gamma_3 < 0$. Because T is tight, some interval (not Δ_1) must go to the leftmost position, and some interval (not Δ_3) must go to the rightmost position. Obviously, there are 3 cases:

	A	A'	$B \& C$
<i>Leftmost</i>	Δ_2	Δ_3	Δ_3
<i>Rightmost</i>	Δ_1	Δ_2	Δ_1

The cases A and A' are mirror images of each other, so we consider only case A of these two.

2.1. Case A. Double rotation in disguise. In this case, Δ_2 goes to the leftmost position and Δ_1 goes to the rightmost position, see Figure 2.1. Then T is a double rotation with $c = \beta_2$ (i.e. the first arc is $\Delta_1 \cup \Delta_2$ and the second one is Δ_3) and $a = -|\Delta_1|$, $b = \gamma_3$.

2.2. Cases B and C. Induction. In this case, Δ_1 goes to the rightmost position and Δ_3 goes to the leftmost position. Because of the symmetry, we can assume without loss of generality that $|\Delta_1| \geq |\Delta_3|$. Consider the two sub-cases: $\gamma_2 < 0$ and $\gamma_2 > 0$.





FIGURE 2.1. Double rotation in disguise.

Sub-Case $\gamma_2 < 0$ [piece B].

Proposition 2.1. *In this case, $\Delta = \Delta_1 \cup \Delta_2$ is regular with the return time ≤ 2 . T_Δ is a tight ITM of three intervals which is a double rotation.*

FIGURE 2.2. Induction to $\Delta_1 \cup \Delta_2$.

Proof. $\gamma_2 < 0$ implies $T\Delta_2 \subset \Delta$, so Δ_2 returns to Δ in one piece after a single iteration of T . On the other hand, Δ_1 is split by the first return map into two pieces, Δ'_1 and Δ''_1 , separated by the point $\beta_2 - \gamma_1$, see Figure 2.2. For the first piece,

$$T\Delta'_1 = T\Delta_1 \cap \Delta \subset \Delta.$$

For the second piece, we have

$$T\Delta''_1 = T\Delta_1 \setminus \Delta = \Delta_3, \quad \text{and} \quad T^2\Delta''_1 = T\Delta_3 \subset \Delta_1 \subset \Delta.$$

Here we used $|\Delta_1| \geq |\Delta_3|$. We have just shown that Δ is regular. T_Δ is a tight ITM of three intervals, $\Delta'_1, \Delta''_1, \Delta_2$. Note that T_Δ sends Δ'_1 to the rightmost position and Δ''_1 to the leftmost one. Thus we have Case A for T_Δ , and T_Δ is a double rotation.

Note that the induction operator $T \mapsto T_\Delta$ is an invertible rational map on B . \square

Remark 2.2. The sub-case $\gamma_2 > 0$ is a generalized Bruin-Troubetzkoy family [4] with one extra degree of freedom, γ_2 .

Sub-Case $\gamma_2 > 0$ [pieces B_i, C_i].

Proposition 2.3. *In this case, $\Delta = \Delta_2 \cup \Delta_3$ is regular, and T_Δ is a tight ITM of three intervals which is a double rotation.*

Proof. Similarly, $\gamma_2 > 0$ implies $T\Delta_2 \subset \Delta$, so Δ_2 returns to Δ in one piece after a single iteration of T . Let us now observe the return of Δ_3 .

We know $T\Delta_3$ is at the leftmost position, which implies $T\Delta_3 \subset \Delta_1$. Thus $T^2\Delta_3 = T\Delta_3 + \gamma_1$. Moreover, for any $n \in \mathbb{N}$ such that $T^n\Delta_3 \subset \Delta_1$, we have $T^n\Delta_3 = T\Delta_3 + (n-1)\gamma_1$. Let n be the maximal n such that $T^2\Delta_3, \dots, T^n\Delta_3 \subset \Delta_1$. There are two possibilities:

- i) $T^{n+1}\Delta_3 \subset \Delta$, or
- ii) $T^{n+1}\Delta_3 \cap \Delta \neq \emptyset, T^{n+1}\Delta_3 \cap \Delta \neq \emptyset$.

Possibility **i** corresponds to $B_i, i = n$. In this case, Δ is obviously regular, and $T_\Delta \in \text{ITM}(2)$, see Figure 2.3. By [1], T_Δ reduces to a (single) rotation.




FIGURE 2.3. Induction to $\Delta_2 \cup \Delta_3$: possibility **i**.

Possibility **ii** corresponds to $C_i, i = n$. In this case, Δ_3 is split into two pieces Δ'_3 and Δ''_3 . Δ''_3 returns to Δ after $n+1$ iterations of T and goes to the leftmost position, see Figure 2.4. Δ'_3 returns to Δ after $n+2$ iterations and goes to the rightmost position. Thus we have Case A' for T_Δ , and T_Δ is a double rotation.




FIGURE 2.4. Induction to $\Delta_2 \cup \Delta_3$: possibility **ii**.

Note that the induction operator $T \mapsto T_\Delta$ is an invertible rational map on each B_i or C_i . \square

3. PROOF OF THEOREM 1.1

Note that (apart from a few trivial exceptions) for any $T \in \text{ITM}(3)$ the limit set X is either entirely periodic or transitive. All the infinite type ITMs belong to the latter

case. Also for ITM(3) we can ignore the case (1.1), because in this case the ITM is just a rotation, and cannot be infinite type. Recall that by Lemma 1.3, the properties to be finite or infinite type are preserved by inductions.

Now we invoke Theorem 1.5 by Bruin and Clack [3]. Together with our Theorem 1.6 it implies that the set $\mathcal{S} \cap \text{TITM}(3)$ is a countable union of sets of zero Lebesgue measure. [3] also numerically estimated the Hausdorff dimension of the infinite type parameters to be between 2 and 2.88. Thus,

$$\text{Leb}(\mathcal{S} \cap \text{TITM}(3)) = 0, \quad 2 \leq \dim(\mathcal{S} \cap \text{TITM}(3)) \leq 2.88.$$

By Proposition 1.4, the set $\mathcal{S} \cap \text{ITM}(3) = \mathcal{F}^{-1}(\mathcal{S} \cap \text{TITM}(3))$ has zero Lebesgue measure and Hausdorff dimension between 4 and 4.88. The main theorem is proven.

ACKNOWLEDGEMENTS

The author is grateful to Professor A. Gorodetski for many fruitful discussions, to University of California, Irvine, for hospitality during the initial work on this paper, and to the anonymous referee whose comments helped to improve the paper. The graphical representations of ITMs are inspired by the figures of interval identifications by A. Skripchenko in [11].

REFERENCES

- [1] Michael Boshernitzan and Isaac Kornfeld. Interval translation mappings. *Ergodic Theory Dynam. Systems*, 15(5):821–832, 1995.
- [2] Henk Bruin. Renormalization in a class of interval translation maps of d branches. *Dyn. Syst.*, 22(1):11–24, 2007.
- [3] Henk Bruin and Gregory Clack. Inducing and unique ergodicity of double rotations. *Discrete and Continuous Dynamical Systems - Series A*, 32(12):4133–4147, 2012.
- [4] Henk Bruin and Serge Troubetzkoy. The Gauss map on a class of interval translation mappings. *Israel J. Math.*, 137:125–148, 2003.
- [5] Jérôme Buzzi. Piecewise isometries have zero topological entropy. *Ergodic Theory Dynam. Systems*, 21(5):1371–1377, 2001.
- [6] Jérôme Buzzi and Pascal Hubert. Piecewise monotone maps without periodic points: rigidity, measures and complexity. *Ergodic Theory Dynam. Systems*, 24(2):383–405, 2004.
- [7] Arek Goetz. Sofic subshifts and piecewise isometric systems. *Ergodic Theory Dynam. Systems*, 19(6):1485–1501, 1999.
- [8] Arek Goetz. Dynamics of piecewise isometries. *Illinois J. Math.*, 44(3):465–478, 2000.
- [9] Arek Goetz. Stability of piecewise rotations and affine maps. *Nonlinearity*, 14(2):205–219, 2001.
- [10] Jörg Schmeling and Serge Troubetzkoy. Interval translation mappings. In *Dynamical systems (Luminy-Marseille, 1998)*, pages 291–302. World Sci. Publ., River Edge, NJ, 2000.
- [11] Alexandra Skripchenko. Symmetric interval identification systems of order three. *arXiv:1010.1820*, 2011.
- [12] Hideyuki Suzuki, Shunji Ito, and Kazuyuki Aihara. Double rotations. *Discrete Contin. Dyn. Syst.*, 13(2):515–532, 2005.

DENIS VOLK
KUNGLIGA TEKNISKA HÖGSKOLAN (ROYAL INSTITUTE OF TECHNOLOGY)
INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS, RUSSIAN ACADEMY OF SCIENCES

E-mail address: `dvolk@kth.se`